





Fax Completed Form To 1.833.404.2392

Prescriber Help Desk 1.833.587.2012

Request for Prior Authorization Dupilumab (Dupixent)

Online <u>covermymeds.com/main/</u> prior-authorization-forms/

(PLEASE PRINT - ACCURACY IS IMPORTANT)

IA Medicaid Member ID #	Patient name	DOB	
Patient address			
Provider NPI	Prescriber name	Phone	
Prescriber address		Fax	
Pharmacy name	Address	Phone	
Prescriber must complete all information above. It must be legible, correct, and complete or form will be returned.			
Pharmacy NPI	Pharmacy fax	NDC	
	-		

Prior authorization is required for Dupixent (dupilumab). Payment for non-preferred agents will be considered when there is documentation of a previous trial and therapy failure with a preferred agent. Payment will be considered when patient has an FDA approved or compendia indication for the requested drug under the following conditions:

- I) Request adheres to all FDA approved labeling for requested drug and indication including age, dosing, contraindications, warnings and precautions, drug interactions, and use in specific populations; and
- 2) Patient's current weight in kilograms (kg) is provided; and
- 3) Patient has a diagnosis of moderate-to-severe atopic dermatitis; and
 - a. Patient has failed to respond to good skin care and regular use of emollients; and
 - b. Patient has documentation of an adequate trial and therapy failure with one preferred medium to high potency topical corticosteroid for a minimum of 2 consecutive weeks; and
 - c. Patient has documentation of a previous trial and therapy failure with a topical immunomodulator for a minimum of 4 weeks; and
 - d. Patient will continue with skin care regimen and regular use of emollients; or
- 4) Patient has a diagnosis of moderate to severe asthma with an eosinophilic phenotype (with a pretreatment eosinophil count ≥ 150 cells/mcL within the previous 6 weeks) OR with oral corticosteroid dependent asthma; and
 - a. Has a pretreatment forced expiratory volume in 1 second (FEV₁) ≤ 80% predicted in adults; < 90% predicted in adolescents 12 to 17 years of age; and < 95% predicted in children 6 to 11 years of age; and
 - b. Symptoms are inadequately controlled with documentation of current treatment with a high-dose inhaled corticosteroid (ICS) given in combination with a controller medication (e.g. long acting beta₂ agonist [LABA], leukotriene receptor antagonist [LTRA], oral theophylline) for a minimum of 3 consecutive months. Patient must be compliant with therapy, based on pharmacy claims; and
 - c. Patient must have one of the following, in addition to the regular maintenance medications defined above:
 - i. One (1) or more exacerbations in the previous year, or
 - ii. Require daily oral corticosteroids for at least 3 days; or
- 5) Patient has a diagnosis of inadequately controlled chronic rhinosinusitis with nasal polyposis (CRSwNP); and
 - a. Documentation dupilumab will be used as an add-on maintenance treatment; and
 - b. Documentation of an adequate trial and therapy failure with at least one preferred medication from each of the following categories:
 - i. Nasal corticosteroid spray; and
 - ii. Oral corticosteroid; or
- 6) Patient has a diagnosis of eosinophilic esophagitis (EoE); and
 - a. Patient has ≥ 15 intraepithelial eosinophils per high-power field (eos/hpf) as confirmed by endoscopic esophageal biopsy (attach results); and
 - b. Patient has signs and symptoms of esophageal dysfunction (e.g., dysphagia, food impaction, food refusal, abdominal pain. heartburn regurgitation, chest pain and/or, odynophagia); and
 - c. Documentation of previous trials and therapy failures with all of the following:
 - i. High dose proton pump inhibitor (PPI) for at least 8 weeks; and
 - ii. Swallowed topical corticosteroid (e.g., fluticasone propionate, oral budesonide suspension); and
 - iii. Dietary therapy; or



ြာ Iowa Health Link 🔨

Hawki

Request for Prior Authorization Dupilumab (Dupixent)

Fax Completed Form To 1.833.404.2392 Prescriber Help Desk 1.833.587.2012

Online

covermymeds.com/main/ prior-authorization-forms/

(PLEASE PRINT - ACCURACY IS IMPORTANT)

- 7) Patient has a diagnosis of moderate to severe prurigo nodularis (PN); and
 - a. Patient has experienced severe to very severe pruritis, as demonstrated by a current Worst Itch-Numeric Rating Scale (WI-NRS) ≥ 7; and
 - b. Patient has \geq 20 nodular lesions (attach documentation); and
 - c. Documentation of a previous trial and therapy failure with a high or super high potency topical corticosteroid for at least 14 consecutive days; or
- 8) Patient has a diagnosis of chronic obstructive pulmonary disease (COPD) and an eosinophilic phenotype; and
 - a. Patient has moderate to severe airflow limitation, measured within the past 12 months, as evidenced by both of the following:
 - i. FEV1/FVC ratio < 0.7, and

i.

- ii. FEV1 % predicted between 30% to79%; and
- b. Patient has a minimum blood eosinophil count of 300 cells/mcL, measured within the past 12 months; and
- c. Patient has documentation of maximal inhaled therapy for 3 or more months and an inadequate response to:
 - Triple therapy with all of the following treatments:
 - 1. Long-acting muscarinic antagonist/anticholinergic (LAMA); and
 - 2. Long-acting beta agonist (LABA); and
 - 3. Inhaled corticosteroid (ICS); or
 - ii. Double therapy with all of the following if ICS is contraindicated
 - 1. LABA; and
 - 2. LAMA; and
- d. Patient has history of at least 2 moderate or 1 severe exacerbation(s) in the previous 12 months despite receiving maximal triple therapy or double therapy (defined above). Moderate exacerbation is defined as patient required treatment with systemic corticosteroids and/or antibiotics and severe exacerbation is defined as hospitalization or observation for over 24 hours in an emergency department or urgent care facility; and
- e. Patient will continue to receive maintenance therapy (as documented above) concomitantly with dupilumab; and
- 9) Dose does not exceed the FDA approved dosing for indication.

If criteria for coverage are met, initial authorizations will be given for 6 months for all the above indications, except COPD, which will receive an initial authorization of 12 months to assess the response to treatment. Requests for continuation of therapy will require documentation of a positive response to therapy.

The required trials may be overridden when documented evidence is provided that use of these agents would be medically contraindicated.

Preferred

Dupixent			
Strength	Usage Instructions	Quantity	Day's Supply
Diagnosis:			
Patient's current weight in k	g:Date o	btained:	
Moderate-to-Severe Atop	ic Dermatitis		
Did patient fail to respond to	good skin care and regular use of e	mollients?	
Yes No If yes, provi	de documentation below:		
Provide skin care regimen, inc	luding name and dates of emollient use	:	

	iowa total care.	டு Iowa Hea	lth Link Iowa HHS	V Hawki	Fax Completed Form To 1.833.404.2392
	•	uest for Prior Au upilumab (Du		1	Prescriber Help Desk 1.833.587.2012 Online
	(PLEASE P	RINT – ACCURAC	Y IS IMPOR	TANT)	<u>covermymeds.com/main/</u> prior-authorization-forms/
Will patient continue skin c	are regimen and	regular use of em	nollients?	Yes No	
Preferred medium to high p	otency topical c	orticosteroid trial	:		
Drug name & dose:			Trial dates		
Failure reason:					
Topical immunomodulator	trial:				
Drug name & dose:			Trial dates	:	
Failure reason:					
Moderate-to-Severe Ast	hma with an Eos	inophilic Phenoty	vpe		
Does patient have pretreatr		count ≥ 150 cells/i	mcL within t	he previous 6 wee	ks?
Does patient have oral corti Yes No Provide pretreatment FEV ₁					
Document current treatmen	it with a high-do	se ICS given in co	mbination w	rith a controller me	dication:
High-Dose ICS Trial:					
Drug name & dose:			Trial dates		
Failure reason:					
Controller Medication Trial:					
Drug name & dose:			Trial dates		
Failure reason:					
Does patient have one of th	e following?				
One (1) or more exacerbation	is in the previous	year? 🗌 Yes 🗌] No		
Require daily oral corticoster	oids for at least 3	days? 🗌 Yes 🗌] No		
Inadequately controlled	chronic rhinosi	nusitis with nasal	polyposis (0	CRSwNP)	
Will dupliumab be used as	an add-on maint	enance treatment	?		
Yes (document concomitaNo	ant maintenance t	reatment): Drug na	ime & dose: _		
Document adequate trial an following categories:	ld therapy failure	e with at least one	preferred m	edication from eac	ch of the
Nasal Corticosteroid Spray	Trial:				
Drug name & dose:			Trial dates	:	
Failure reason:					







Request for Prior Authorization Dupilumab (Dupixent)

1.833.587.2012 Online covermymeds.com/main/ prior-authorization-forms/

(PLEASE F	PRINT –	ACCURACY	IS IMP	ORTANT)
(•••••

Oral Corticosteroid Trial:		prior-authorization-forms/
Drug name & dose:	Trial dates:	
Failure reason:		
Eosinophilic Esophagitis (Eo	Е)	
Does patient have ≥ 15 intraepithe endoscopic esophageal biopsy?□ Yes (attach results)□ No	elial eosinophils per high-power field (eos/hpf) confirme	d by
	nptoms of esophageal dysfunction?	
Document previous trials and the	erapy failures with all of the following:	
High Dose PPI :		
Drug name & dose:	Trial dates:	
Failure reason:		
Swallowed topical corticosteroid:	:	
Drug name & dose:	Trial dates:	
Failure reason:		
Dietary Therapy:		
Dietary Plan:	Trial dates:	
Failure reason:		
Moderate to Severe Prurigo N	lodularis (PN)	
Worst Itch-Numeric Rating Scale ((WI-NRS) response: Date	e obtained:
Does patient have ≥ 20 nodular le	esions? 🗌 Yes (provide documentation) 🔲 No	
Preferred high or super high pote	ency topical corticosteroid trial:	
Drug name & dose:	Trial dates:	
Failure reason:		
Chronic Obstructive Pulmona	ary Disease (COPD) and an eosinophilic phenotype:	
Provide all of the following inform	nation:	
FEV1/FVC ratio:	Date obtained:	
FEV1 % predicted:	Date obtained:	







Fax Completed Form To 1.833.404.2392 Prescriber Help Desk 1.833.587.2012

Online covermymeds.com/main/

prior-authorization-forms/

Request for Prior Authorization Dupilumab (Dupixent)

(PLEASE PRINT – ACCURACY IS IMPORTANT)

Blood eosinophil count:	Date obtained:
Trial information:	
LABA Trial:	
Name/dose:	Trial dates:
Failure reason/medical contraindication:	
LAMA Trial:	
Name/dose:	Trial dates:
Failure reason/medical contraindication:	
ICS Trial:	
Name/dose:	Trial dates:
Failure reason/medical contraindication:	
Document exacerbations:	
Moderate:	
Date:	_ Treatment needed:
Date:	_ Treatment needed:
Severe:	
Date:	Place of care:
Will patient continue to receive main	tenance therapy concomitantly with dupilumab? Yes No
Renewal requests:	
Document positive response to therapy:_	
Attach lab results and other documen	tation as necessary.

Prescriber signature (Must match prescriber listed above.) Date of submission

IMPORTANT NOTE: In evaluating requests for prior authorization the consultant will consider the treatment from the standpoint of medical necessity only. If approval of this request is granted, this does not indicate that the member continues to be eligible for Medicaid. It is the responsibility of the provider who initiates the request for prior authorization to establish by inspection of the member's Medicaid eligibility card and, if necessary by contact with the county Department of Health and Human Services, that the member continues to be eligible for Medicaid.