





Fax Completed Form To 1.833.404.2392

Prescriber Help Desk 1.833.587.2012

Online covermymeds.com/main/prior-

Request for Prior Authorization Dupilumab (Dupixent)

(PLEASE PRINT - ACCURACY IS IMPORTANT)

	(PLEASE PRINT – ACCURACY IS IMPOR	(IANI) <u>authorization-forms/</u>
IA Medicaid Member ID #	Patient name	DOB
Patient address		
Provider NPI	Prescriber name	Phone
Prescriber address		Fax
Pharmacy name	Address	Phone
Prescriber must complete all informa	ition above. It must be legible, correct, and comp	lete or form will be returned.
Pharmacy NPI	Pharmacy fax N	DC

Prior authorization is required for Dupixent (dupilumab). Payment for non-preferred agents will be considered when there is documentation of a previous trial and therapy failure with a preferred agent. Payment will be considered when patient has an FDA approved or compendia indication for the requested drug under the following conditions:

- Request adheres to all FDA approved labeling for requested drug and indication including age, dosing, contraindications, warnings and precautions, drug interactions, and use in specific populations; and
- 2) Patient's current weight in kilograms (kg) is provided; and
- Patient has a diagnosis of moderate-to-severe atopic dermatitis; and
 - Is prescribed by or in consultation with a dermatologist, allergist, or immunologist; and
 - Patient has failed to respond to good skin care and regular use of emollients; and b.
 - Patient has documentation of an adequate trial and therapy failure with one preferred medium to high potency topical corticosteroid for a minimum of 2 consecutive weeks; and
 - Patient has documentation of a previous trial and therapy failure with a topical immunomodulator for a minimum of 4 weeks; and
 - Patient has documentation of a previous trial and therapy failure with cyclosporine or azathioprine; and
 - Patient will continue with skin care regimen and regular use of emollients; or
- Patient has a diagnosis of moderate to severe asthma with an eosinophilic phenotype (with a pretreatment eosinophil count ≥ 150 cells/mcL within the previous 6 weeks) OR with oral corticosteroid dependent asthma; and
 - Is prescribed by or in consultation with an allergist, immunologist, or pulmonologist; and
 - Has a pretreatment forced expiratory volume in I second (FEV_I) \leq 80% predicted; and b.
 - Symptoms are inadequately controlled with documentation of current treatment with a high-dose inhaled corticosteroid (ICS) given in combination with a controller medication (e.g. long acting beta2 agonist [LABA], leukotriene receptor antagonist [LTRA], oral theophylline) for a minimum of 3 consecutive months. Patient must be compliant with therapy, based on pharmacy claims; and
 - Patient must have one of the following, in addition to the regular maintenance medications defined above:
 - Two (2) or more exacerbations in the previous year, or
 - Require daily oral corticosteroids for at least 3 days; or
- Patient has a diagnosis of inadequately controlled chronic rhinosinusitis with nasal polyposis (CRSwNP); and
 - Documentation dupilumab will be used as an add-on maintenance treatment; and
 - Documentation of an adequate trial and therapy failure with at least one preferred medication from each of the following categories:
 - Nasal corticosteroid spray; and
 - Oral corticosteroid; or
- Patient has a diagnosis of eosinophilic esophagitis (EoE); and
 - Is prescribed by, or in consultation with, and allergist, gastroenterologist, or immunologist; and
 - Patient has ≥ 15 intraepithelial eosinophils per high-power field (eos/hpf) as confirmed by endoscopic esophageal biopsy (attach results); and
 - Patient has signs and symptoms of esophageal dysfunction (e.g., dysphagia, food impaction, food refusal, abdominal pain. heartburn regurgitation, chest pain and/or, odynophagia); and
 - Documentation of previous trials and therapy failures with all of the following:

Rev. 6/24 Page I of 5

(PLEASE PRINT - ACCURACY IS IMPORTANT)

- i. High dose proton pump inhibitor (PPI) for at least 8 weeks; and
- ii. Swallowed topical corticosteroid (e.g., fluticasone propionate, oral budesonide suspension); and
- iii. Dietary therapy; or
- 7) Patient has a diagnosis of moderate to severe prurigo nodularis (PN); and
 - a. Is prescribed by, or in consultation with an allergist, immunologist, or dermatologist; and
 - b. Patient has experienced severe to very severe pruritis, as demonstrated by a current Worst Itch-Numeric Rating Scale (WI-NRS) ≥ 7; and
 - c. Patient has ≥ 20 nodular lesions (attach documentation); and
 - d. Documentation of a previous trial and therapy failure with a high or super high potency topical corticosteroid for at least 14 consecutive days; and
- 8) Dose does not exceed the FDA approved dosing for indication.

If criteria for coverage are met, initial authorizations will be given for 6 months to assess the response to treatment. Requests for continuation of therapy will require documentation of a positive response to therapy.

The required trials may be overridden when documented evidence is provided that use of these agents would be medically contraindicated.

<u>Preferred</u>			
Dupixent			
Strength	Usage Instructions	Quantity	Day's Supply
Diagnosis:			
Patient's current weight in l	kg:Date ob	otained:	
☐ Moderate-to-Severe Ato	opic Dermatitis		
Is prescriber a dermatologis	st, allergist, or immunologist?		
Yes specialty:			
☐ No If no, note consultation	on with dermatologist, allergist, or immunolog	gist:	
Consultation date:	Physician name, specialty & phone:	:	
Did patient fail to respond t	o good skin care and regular use of emo	ollients?	
Yes No If yes, prov	ide documentation below:		
Provide skin care regimen, inclu	ding name and dates of emollient use:		
Will patient continue skin ca	are regimen and regular use of emollier	nts? Yes No	
Preferred medium to high p	ootency topical corticosteroid trial:		
Drug name & dose:	Tri	ial dates:	
Failure reason:			
Topical immunomodulator	trial:		
Drug name & dose:	Tri	ial dates:	
Failure reason:			

Rev. 6/24 Page 2 of 5

(PLEASE PRINT – ACCURACY IS IMPORTANT)

Drug name & dose:Trial dates:Trial dates:				
Failure reason:				
Medical or contraindication reason to override trial requirements:				
■ Moderate-to-Severe Asthma with an Eosinophilic Phenotype				
Does patient have pretreatment eosinophil count ≥ 150 cells/mcL within the previous 6 weeks? Yes (attach results) No No No Yes No				
Is prescriber an allergist, immunologist, or pulmonologist?				
Yes specialty:				
No If no, note consultation with allergist, immunologist, or pulmonologist:				
Consultation date: Physician name, specialty & phone:				
Does patient have a pretreatment FEV₁ ≤ 80% predicted? ☐ Yes (attach results) ☐ No				
Document current treatment with a high-dose ICS given in combination with a controller medication:				
High-Dose ICS Trial:				
Drug name & dose:Trial dates:				
Failure reason:				
Controller Medication Trial:				
Drug name & dose:Trial dates:				
Failure reason:				
Does patient have one of the following?				
Two (2) or more exacerbations in the previous year? Yes No				
Require daily oral corticosteroids for at least 3 days? Yes No				
☐ Inadequately controlled chronic rhinosinusitis with nasal polyposis (CRSwNP)				
Will dupliumab be used as an add-on maintenance treatment?				
Yes (document concomitant maintenance treatment): Drug name & dose:				
□ No				
Document adequate trial and therapy failure with at least one preferred medication from each of the following categories:				
Nasal Corticosteroid Spray Trial:				
Drug name & dose:Trial dates:				
Failure reason:				

Rev. 6/24 Page 3 of 5

(PLEASE PRINT – ACCURACY IS IMPORTANT)

Oral Corticosteroid Trial:				
Drug name & dose:	Trial dates:			
Failure reason:				
☐ Eosinophilic Esophagitis (Eo	DE)			
Is prescriber an allergist, immu	nologist, or gastroenterologist?			
Yes specialty:				
☐ No If no, note consultation w	No If no, note consultation with allergist, immunologist, or gastroenterologist:			
Consultation date:	Physician name, specialty & phone:			
Does patient have ≥ 15 intraep esophageal biopsy?	thelial eosinophils per high-power field (eos/hpf) confirmed by endoscopic			
Yes (attach results) No				
	mptoms of esophageal dysfunction?			
Document previous trials and therapy failures with all of the following:				
High Dose PPI:				
Drug name & dose:	Trial dates:			
Failure reason:				
Swallowed topical corticostero	id:			
Drug name & dose:	Trial dates:			
Failure reason:				
Dietary Therapy:				
Dietary Plan:	Trial dates:			
Failure reason:				
☐ Moderate to Severe Prurig	o Nodularis (PN)			
Is prescriber an allergist, immu	nologist, or dermatologist?			
Yes specialty:				
☐ No If no, note consultation w	rith allergist, immunologist, or dermatologist:			
Consultation date:	Physician name, specialty & phone:			
Worst Itch-Numeric Rating Sca	ale (WI-NRS) response: Date obtained:			
Does patient have \geq 20 nodular	lesions? Yes (provide documentation) No			

Rev. 6/24 Page 4 of 5

(PLEASE PRINT - ACCURACY IS IMPORTANT)

Preferred high or super high potency topical corticosteroid trial:

Drug name & dose:	Trial dates:	
Failure reason:		
Renewal requests:		
Document positive response to therapy:		
And the first of the second		
Attach lab results and other documentation as necessary.		
Prescriber signature (Must match prescriber listed above.)	Date of submission	

IMPORTANT NOTE: In evaluating requests for prior authorization the consultant will consider the treatment from the standpoint of medical necessity only. If approval of this request is granted, this does not indicate that the member continues to be eligible for Medicaid. It is the responsibility of the provider who initiates the request for prior authorization to establish by inspection of the member's Medicaid eligibility card and, if necessary by contact with the county Department of Health and Human Services, that the member continues to be eligible for Medicaid.

Rev. 6/24 Page 5 of 5