





Request for Prior Authorization SHORT ACTING OPIOIDS

(PLEASE PRINT - ACCURACY IS IMPORTANT)

Fax Completed Form To 1.833.404.2392

Prescriber Help Desk 1.833.587.2012

Online

covermymeds.com/main/ prior-authorization-forms/

IA Medicaid Member I	D#	Patient name			DOB		
Patient address							
B : I NE		Т					
Provider NPI	1 1 1	Prescriber name			Phone		
Prescriber address					Fax		
		1					
Pharmacy name		Address			Phone		
	plete all infori	mation above. It must be legible,			rm will be	returned.	
Pharmacy NPI		Pharmacy fax	ND	С			
use of controlled sul of a short-acting of addiction, abuse and adverse effects and prescriber must do discussed with the p plan to taper the be be given for 3 mos experienced improve controlled substant appropriate for this following: a. the risk Documentation as to is provided, if appro- these agents and/or	ostances on the bioid is appropriate. The recomment the patient; and be needed to b	ngredient only) at therapeutic he lowa Prescription Monitoric priate for this member base to requesting prior authorizerse effects of opioids; and 7 following: a. The risks of using Documentation as to why consist is provided, if appropriate. If conal approvals will be considered nontrol and level of functions own PMP website and has control and benzodiazepines concern use is medically necessal equired trials may be overrided to logic therapies would be measured.	ng Program (PMP ed on review of I ation; and 6) Pation) For patients taling opioids and be neurrent use is moriteria for coveragered if the following; and 2) Prescribetermined contint benzodiazeping oncurrently has be ry is provided; and den when docume) website PMP and PMP	te and has d the pat been infor neurrent tepines co necessar net, an ini teria are us reviewe se of a s orescriber ussed wit an to tape	s determing tient's risk remed of the benzodiaze oncurrently is provide tial authored the pation of the pation of the pation of the pation of the benzelon.	ned that us of for opioine commo epines, the ly has been led; and c. Arization wife a popioid in the lent's use of the lent, and le ent, a
Preferred (*Please reficomplete list of prefer Acetaminophen/Codeine Hydrocodone/APAP Hydromorphone Tab Morphine Sulfate Tab Oxycodone Cap/Tab Oxycodone /APAP (5/325) Tramadol 50mg	red alternative	Butalbital/APAP/Caff/ Butalbital/ASA/Caff/ Combunox	Codeine (5/300, 7.5/300, 10/30	C C C P R	lucynta Dxymorphon Dxycodone/A rimlev oxicodone ramadol 25n	APAP (7.5/32	5, 10/325)
!	Strength	Dosage Instructions	Quantity	D	ays Supply	У	
	J	3	/		, ,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	•	
Diagnosis:							

Rev. 6/24 Page I of 3

Request for Prior Authorization SHORT ACTING OPIOIDS

(PLEASE PRINT - ACCURACY IS IMPORTANT)

Document non-pharmacologic therapies (such as physical therapy, weight loss, alternative therapies such as manipulation, massage, and acupuncture, or psychological therapies such as cognitive behavior therapy [CBT], etc,)

Trial Dates:	Non-Pharmacological Treatment Trial#I			
Document 2 nonopioid pharmacologic therapies (acetaminophen or NSAIDs) Nonopioid Pharmacologic Trial #1: Name/Dose:	Trial Dates:Failure reason _	· · · · · · · · · · · · · · · · · · ·		
Trial Dates:Failure reason	Non Pharmacological Treatment Trial #2			
Document 2 nonopioid pharmacologic therapies (acetaminophen or NSAIDs) Nonopioid Pharmacologic Trial #1: Name/Dose:				
Nonopioid Pharmacologic Trial #1: Name/Dose: Trial Dates: Trial Dates: Trial Dates: Trial Dates:	, and e reason _		· · · · · · · · · · · · · · · · · · ·	
Nonopioid Pharmacologic Trial #2: Name/Dose:	Document 2 nonopioid pharmacologic therap	vies (acetaminophen or NSAIDs)		
Nonopioid Pharmacologic Trial #2: Name/Dose:	Nonopioid Pharmacologic Trial #1: Name/Dose:		Trial Dates:	
Document trials with three preferred chemically distinct short acting opioids Preferred Trial I: Drug Name	Failure reason			
Document trials with three preferred chemically distinct short acting opioids Preferred Trial 1: Drug Name	Nonopioid Pharmacologic Trial #2: Name/Dose:		Trial Dates:	
Preferred Trial 1: Drug Name Strength Dosage Instructions Trial start date: Trial end date: Preferred Trial 2: Drug Name Strength Dosage Instructions Trial start date: Trial end date: Preferred Trial 2: Drug Name Strength Dosage Instructions Trial start date: Trial end date: Preferred Trial 3: Drug Name Strength Dosage Instructions Trial start date: Trial end date: Preferred Trial 3: Drug Name Strength Dosage Instructions Trial start date: Trial end date: Failure reason: Prescriber review of patient's controlled substances use on the Iowa PMP website: No Yes Date Reviewed: Is short-acting opioid use appropriate for patient based on PMP review and patient's risk for opioid addiction, abus and misuse? No Yes Has patient been informed of the common adverse effects (constipation, dry mouth, nausea, vomiting, drowsiness confusion, tolerance, physical dependence, and withdrawal symptoms when stopping opioids) and serious adverse effects (potentially fatal overdose and development of a potentially serious opioid use disorder) of opioids? No Yes Patients taking concurrent benzodiazepines:				
Preferred Trial 1: Drug Name				
Trial start date:	·	,		
Failure reason: Preferred Trial 2: Drug Name	·	_	_	
Preferred Trial 2: Drug Name Strength Dosage Instructions Trial start date: Trial end date: Failure reason: Preferred Trial 3: Drug Name Strength Dosage Instructions Trial start date: Trial end date: Failure reason: Prescriber review of patient's controlled substances use on the Iowa PMP website: No Yes Date Reviewed: Is short-acting opioid use appropriate for patient based on PMP review and patient's risk for opioid addiction, abuse and misuse? No Yes Has patient been informed of the common adverse effects (constipation, dry mouth, nausea, vomiting, drowsiness confusion, tolerance, physical dependence, and withdrawal symptoms when stopping opioids) and serious adverse effects (potentially fatal overdose and development of a potentially serious opioid use disorder) of opioids? No Yes Patients taking concurrent benzodiazepines:				
Trial start date:Trial end date:	Failure reason:			
Preferred Trial 3: Drug Name Strength Dosage Instructions Trial start date: Trial end date: Failure reason: Prescriber review of patient's controlled substances use on the Iowa PMP website: No Yes Date Reviewed: Is short-acting opioid use appropriate for patient based on PMP review and patient's risk for opioid addiction, abustand misuse? No Yes Has patient been informed of the common adverse effects (constipation, dry mouth, nausea, vomiting, drowsiness confusion, tolerance, physical dependence, and withdrawal symptoms when stopping opioids) and serious adverse effects (potentially fatal overdose and development of a potentially serious opioid use disorder) of opioids? No Yes Patients taking concurrent benzodiazepines:	Preferred Trial 2: Drug Name	Strength	Dosage Instructions	
Preferred Trial 3: Drug Name Strength Dosage Instructions Trial start date: Trial end date: Failure reason: Failure review of patient's controlled substances use on the Iowa PMP website: No Yes Date Reviewed: Is short-acting opioid use appropriate for patient based on PMP review and patient's risk for opioid addiction, abuse and misuse? No Yes Has patient been informed of the common adverse effects (constipation, dry mouth, nausea, vomiting, drowsiness confusion, tolerance, physical dependence, and withdrawal symptoms when stopping opioids) and serious adverse effects (potentially fatal overdose and development of a potentially serious opioid use disorder) of opioids? No Yes Patients taking concurrent benzodiazepines:	Trial start date:Trial end date	ə: <u> </u>	<u></u>	
Trial start date:Trial end date: Failure reason:	Failure reason:			
Prescriber review of patient's controlled substances use on the lowa PMP website: No Yes Date Reviewed: Is short-acting opioid use appropriate for patient based on PMP review and patient's risk for opioid addiction, abuse and misuse? No Yes Has patient been informed of the common adverse effects (constipation, dry mouth, nausea, vomiting, drowsiness confusion, tolerance, physical dependence, and withdrawal symptoms when stopping opioids) and serious adverse effects (potentially fatal overdose and development of a potentially serious opioid use disorder) of opioids? No Yes Patients taking concurrent benzodiazepines:	Preferred Trial 3: Drug Name	Strength	Dosage Instructions	
Prescriber review of patient's controlled substances use on the Iowa PMP website:	Trial start date:Trial end date	ə: <u> </u>		
Is short-acting opioid use appropriate for patient based on PMP review and patient's risk for opioid addiction, abuse and misuse? No Yes Has patient been informed of the common adverse effects (constipation, dry mouth, nausea, vomiting, drowsiness confusion, tolerance, physical dependence, and withdrawal symptoms when stopping opioids) and serious adverse effects (potentially fatal overdose and development of a potentially serious opioid use disorder) of opioids? No Yes Patients taking concurrent benzodiazepines:	Failure reason:			
and misuse? No Yes Has patient been informed of the common adverse effects (constipation, dry mouth, nausea, vomiting, drowsiness confusion, tolerance, physical dependence, and withdrawal symptoms when stopping opioids) and serious adverse effects (potentially fatal overdose and development of a potentially serious opioid use disorder) of opioids? No Yes Patients taking concurrent benzodiazepines:	Prescriber review of patient's controlled subst	tances use on the Iowa PMP	website: No Yes Date Review	wed:
confusion, tolerance, physical dependence, and withdrawal symptoms when stopping opioids) and serious adverse effects (potentially fatal overdose and development of a potentially serious opioid use disorder) of opioids? No Yes Patients taking concurrent benzodiazepines:			d patient's risk for opioid addiction,	abuse
Patients taking concurrent benzodiazepines:	confusion, tolerance, physical dependence, an	d withdrawal symptoms whe	n stopping opioids) and serious adv	
	☐ No ☐ Yes			
	Patients taking concurrent benzodiazepines:			
Have the risks of using opioids and benzodiazepines concurrently been discussed with the patient?	Have the risks of using opioids and benzodiazepines	concurrently been discussed with	hthe patient?	es es
Medical necessity for concurrent use:	Medical necessity for concurrent use:	· 		

Rev. 6/24 Page 2 of 3

Request for Prior Authorization SHORT ACTING OPIOIDS

(PLEASE PRINT - ACCURACY IS IMPORTANT)

Provide plan to taper the benzodiazepine or medical rationale why not appropriate appropriate plan to taper the benzodiazepine or medical rationale why not appropriate plan to taper the benzodiazepine or medical rationale why not appropriate plan to taper the benzodiazepine or medical rationale why not appropriate plan to taper the benzodiazepine or medical rationale why not appropriate plan to taper the benzodiazepine or medical rationale why not appropriate plan to taper the benzodiazepine or medical rationale why not appropriate plan to taper the benzodiazepine or medical rationale why not appropriate plan to taper the benzodiazepine or medical rationale why not appropriate plan to taper the benzodiazepine or medical rationale why not appropriate plan to tape the benzodiazepine plan tape the benzodi	oriate:						
Renewals							
Has patient experienced improvement in pain control and level of f	unctioning?						
☐ No ☐ Yes (describe)							
Updated prescriber review of patient's controlled substances use of ☐ No ☐ Yes Date Reviewed:	n the Iowa PMP website (s	since initial request):					
Continued use of a short-acting opioid is appropriate for this member?							
☐ No ☐ Yes (describe)							
Patients taking concurrent benzodiazepines:							
Have the risks of using opioids and benzodiazepines concurrently been discus	Have the risks of using opioids and benzodiazepines concurrently been discussed with the patient?						
Medical necessity for concurrent use:							
Provide plan to taper the benzodiazepine or medical rationale why not approp	oriate:						
Other medical conditions to consider Attach lab results and other documentation as necessary.							
Prescriber signature (Must match prescriber listed above.)	Date of submission						

IMPORTANT NOTE: In evaluating requests for prior authorization the consultant will consider the treatment from the standpoint of medical necessity only. If approval of this request is granted, this does not indicate that the member continues to be eligible for Medicaid. It is the responsibility of the provider who initiates the request for prior authorization to establish by inspection of the member's Medicaid eligibility card and, if necessary by contact with the county Department of Health and Human Services, that the member continues to be eligible for Medicaid.

Rev. 6/24 Page 3 of 3